

SPYCRA CONTACT WOUND CARE PROTOCOL DETERSION OF A WOUND IN COMBINATION WITH AN HYDROGEL.



Deep wound with some necrotic tissue



An hydrogel is applied in the wound to enhance wound delectation. Spycra Contact is applied over the hydrogel; an extra absorbent dressing is applied to absorb the excess of fluid.



The secondary absorbent dressing can be changed while Spycra Contact stays in place. Spycra Contact can be lifted to evaluate the healing process.



The soft silicone adhesive of Spycra Contact secures that the surrounding skin as well as the newly formed granulation tissue stay in optimal condition. Dressing changes are a-traumatic and painless.

INSTRUCTIONS FOR USE

1. CLEAN AND DRY THE WOUND AS WELL AS THE SURROUNDING SKIN CAREFULLY.
2. APPLY AN HYDROGEL ON THE WOUND AND CHOOSE A SPYCRA CONTACT LARGE ENOUGH TO COVER THE ENTIRE WOUND AREA. MAKE SURE THAT THE DRESSING OVERLAPS THE WOUND BY 2CM MINIMUM.
3. APPLY AN EXTRA ABSORBENT DRESSING OVER SPYCRA CONTACT IN ORDER TO ABSORB THE EXCESS OF WOUND FLUID.
4. LEAVE SPYCRA CONTACT IN PLACE AS LONG AS THERE ARE NO SIGNS OF INFECTION.

Note: SPYCRA CONTACT can be left in place for up to 7days. If needed, the secondary absorbent dressing can be changed while SPYCRA CONTACT stays in place. SPYCRA CONTACT can be lifted to evaluate the healing process. After inspection, reposition the SPYCRA CONTACT over the wound.